



HUMAN PAPILLOMAVIRUS ASSOCIATED CANCERS

Most Human Papillomavirus (HPV) associated cancers can be **prevented with vaccination**. Nearly all adults will be exposed to HPV in their lifetime. Most will never know they have been infected. Idaho Goal: 80% vaccination rate of Idahoans aged 13-17 by 2025.



HPV-ASSOCIATED CANCERS

Incidence 2017-2021 (rates per 100,000)



CALL TO ACTION

- Protect your child before they are exposed to cancer-causing infections.
- HPV vaccination is recommended for boys and girls starting at age 9.
- Recommend the HPV vaccination along with other immunizations, not as a separate recommendation.
- HPV vaccination is cancer prevention!

To learn more, visit

hpvfreeidaho.org. Free screening and diagnostic testing is offered to low-income, uninsured women by Women's Health Check.

Follow the QR code to learn more and see if you qualify.



This publication has been funded in whole or in part with Federal funds from the National Cancer Institute (NCI) under Contract No. HHSN261201800006I and the Centers for Disease Control and Prevention (CDC) under the following Grants or Cooperative Agreements 1NU58DP007160, NU58DP006314 and DP006084. Its contents are solely the responsibility of the authors and do not necessarily represent the official position of CDC or NCI. January 2024.









