

Colorectal Cancer in Idaho

Colorectal cancer rates are 20% higher in Idaho communities with elevated rates of poverty. Colorectal cancer rates are 25% higher among American Indian Idahoans than among white Idahoans.

DEFEATING COLORECTAL CANCER:

- 45–75: the recommended age for a colorectal cancer screening.
- Early detection saves lives! Colorectal cancer can be prevented or treated if found early.
- Choose your best screening method. You have several screening choices, including simple take-home options.

Rates of colorectal cancer cases have decreased 3.7% between 2003–2022.

1,341

deaths from colorectal cancer among Idahoans from 2019–2023

Let's raise the bar!

3 in 5

eligible Idahoans are NOT up to date on their colorectal cancer screening*



Be here today and tomorrow.



Rates of colorectal cancer deaths have decreased 1.0% between 2004–2023.

* 2022 BRFSS



**WOMEN'S
HEALTH CHECK**




IDAHO DEPARTMENT OF
HEALTH & WELFARE



**Cancer Data
Registry of Idaho**



**COMPREHENSIVE
CANCER CONTROL
OF IDAHO**



**Follow the QR code
to learn more.**

This publication has been funded in whole or in part with Federal funds from the National Cancer Institute (NCI) under Contract No. HHSN261201800006I and the Centers for Disease Control and Prevention (CDC) under the following Grants or Cooperative Agreements NU58DP007160, NU58DP006314 and DP006084. Its contents are solely the responsibility of the authors and do not necessarily represent the official position of CDC or NCI. August 2025